

**g-FACTOR OF THE keV 5/2⁻ STATE IN ¹⁹⁷Pt MEASURED
BY THE TDPAC METHOD**

R. N. Saxena and J. C. Soares

**PUBLICAÇÃO IPEN 38
IPEN - Pub 38**

SETEMBRO/1981

CONSELHO DELIBERATIVO

MEMBROS

Prof. Dr. Luiz Cintra do Prado – Presidente

Dr. Edgardo Azevedo Soares Júnior – Vice-Presidente

CONSELHEIROS

Dr. Hécio Modesto da Costa

Dr. Ivano Humbert Marchesi

Prof. Dr. Waldyr Muniz Oliva

Prof. Dr. José Augusto Martins

REPRESENTANTES

Dr. Jacob Charcot Pereira Rios

Dr. Samuel Angarita Ferreira da Silva

SUPERINTENDENTE

Hernani Augusto Lopes de Amorim

**g-FACTOR OF THE keV 5/2⁻ STATE IN ¹⁹⁷Pt MEASURED
BY THE TDPAC METHOD**

R. N. Saxena and J. C. Soares

**CENTRO DE OPERAÇÃO E UTILIZAÇÃO DO REATOR DE PESQUISAS – COURP
ÁREA DE FÍSICA NUCLEAR**

**INSTITUTO DE PESQUISAS ENERGÉTICAS E NUCLEARES
SÃO PAULO – BRASIL**

SÉRIE PUBLICAÇÃO IPEN

INIS Categories and Descriptors

A34

DIFFERENTIAL PAC: Platinum 197

PLATINUM 197: Differential PAC

GAMMA CASCADES: Platinum 197

PLATINUM 197: Gamma cascades

COURP - AFN

Received in September 1980.

Approved for publication in July 1981.

Writing, orthography, concepts and final revision are of exclusive responsibility of the Authors.

g-FACTOR OF THE keV 5/2⁻ STATE IN ¹⁹⁷Pt MEASURED BY THE TDPAC METHOD¹

R. N. Saxena² and J. C. Soares³

ABSTRACT

The g-factor of the 53 keV state in ¹⁹⁷Pt has been measured using the gamma-gamma time differential perturbed angular correlation (TDPAC) method in an external magnetic field of 25.1 kG. The measurements were performed by utilizing the 346 – 53 keV gamma cascade in the decay of 95.4 min ¹⁹⁷Pt. The value of the g-factor was obtained to be $+0.335 \pm 0.010$. This result is compared with the g-factors of similar states in ¹⁹⁵Pt and ^{197,199}Hg.

INTRODUCTION

The platinum isotopes are situated in a most interesting transitional region where the stable shape of the nucleus changes suddenly from a sphere to a highly deformed spheroid. These nuclei therefore offer an excellent opportunity to observe changes between the spherical and spheroidal shapes. Although considerable experimental data on the low-lying levels in odd-A Pt isotopes have been reported, our understanding of the structure of these nuclei is still unsatisfactory. The $P_{1/2,3/2}$, $f_{5/2}$ and $i_{13/2}$ single neutron states in odd-A Pt isotopes are expected to be fairly low in energy⁽⁷⁾, however they have not been identified conclusively except in the case of ¹⁹⁵Pt. Several theoretical calculations have been carried out in the past to explain the observed properties of these nuclei. In the case of ¹⁹⁵Pt Gal² made calculations using the core-excitation model where the energies of the levels, the multipolarities of the transitions and the transition rates between the ground state 1/2⁻ and the two low lying doublets 3/2⁻ and 5/2⁻ are reproduced rather well. However the agreement with the measured magnetic moments of these states was poor with the exception of the first 5/2⁻ state at 130 keV⁽¹²⁾.

The present measurement of the g-factor of the 53 keV 5/2⁻ state in ¹⁹⁷Pt was undertaken with the two fold interest: a) to provide an experimental value of the magnetic moment to further elucidate the nature of this 5/2⁻ state in terms of the collective (core-excitation model) or the quasi particle model; b) to demonstrate the possibility of using the 346-53 keV gamma-cascade in ¹⁹⁷Pt in TDPAC studies. To our knowledge so far all the magnetic hyperfine interaction studies on platinum have been performed using the ¹⁹⁵Pt as the probe nucleus. The 95.4 min 13/2⁺ state of ¹⁹⁷Pt almost exclusively depopulates by the cascade with 346 keV and 53 keV gamma-rays and the large theoretical angular correlation coefficient, $A_{22}^{theo} = 0.2207$, of this cascade makes it quite attractive for the time differential investigation of hyperfine interactions. The measured half life of the 53 keV level $T_{1/2} = 16.58 \pm 0.17$ ns⁽¹⁰⁾ also seems to be quite suitable for this purpose. However it is essential to know the precise value of the nuclear moment of the state before it can be used in the TDPAC studies. We have measured the g-factor of the 53 keV state in ¹⁹⁷Pt using the gamma-gamma TDPAC method in an external magnetic field of 25.1 kG. The measurements were performed by utilizing the constant angle reverse field method and the 346-53 keV gamma cascade in the decay of 95.4 min ^{197m}Pt.

(1) Partial financial support for this work provided by CNEN, Brasil and INIC, Portugal.

(2) Instituto de Pesquisas Energéticas e Nucleares, São Paulo, SP, Brasil.

(3) Centro de Física Nuclear, Universidade de Lisboa, Lisboa, Portugal.

EXPERIMENTAL

The radioactive sources of 95.4 min ^{197m}Pt were produced by the neutron irradiation of thin platinum foils ($\approx 10\text{mg/cm}^2$) containing $\approx 98\%$ ^{196}Pt . The neutron irradiations were carried out in the IEA-R1 reactor with the neutron flux of $\approx 10^{13}$ n/cm 2 .s during 10 minutes. Since repeated irradiations were necessary for the experiment, a number of foils were prepared and any given foil was again irradiated only after a cooling period of 4-5 days in order to reduce the contributions from the decay of 18 h ^{197}Pt . Only other impurity in the sample was that of 30 min ^{199}Pt present in small quantity. A total of 80 irradiations were carried out for the entire experiment.

For our time differential experiment we utilized two 2" x 2" NaI(Tl) detectors coupled to RCA 8850 and RCA 8575 phototubes through the 30 cm lucite light guides to detect the 53 keV and 346 keV gamma radiations respectively. A conventional fast slow coincidence system utilizing the differential discriminators and a time to pulse height converter in connection with a multichannel analyser was used for recording the delayed coincidence spectrum. The timing resolution of the equipment was determined by using the 343 keV gamma - 53 keV X-ray cascade in the decay of ^{175}Hf . Typical time resolution of the set up was 5.5 ns FWHM. The use of the long light guides was probably responsible for this rather large time resolution. A FWHM of 2.9 ns was obtained in our experiment⁽¹⁰⁾ for the lifetime measurement of the 53 keV state in ^{197}Pt where no light guides were used. The magnetic field of 25.1 kG was supplied by an electromagnet. The delayed coincidence spectra for each direction of the magnetic field were stored in two different subgroups of the multichannel analyser memory. The field direction was changed every 10 min with the detectors maintained at 135° . In this manner each source was measured for 180 min before replacing it with a fresh source.




Figure 1 - Spin precession of the magnetic moment of the 53 keV state in ^{197}Pt in the external magnetic field of 25.1 kG.

RESULTS AND DISCUSSION

The asymmetry ratio $T(t) = (N\uparrow - N\downarrow)/(N\uparrow + N\downarrow)$ calculated from the measured time spectra are plotted in Figure 1. The solid curve is the least square fit of the experimental data to the function $R(t) = A \sin 2\omega_L t$. The data corresponding to the initial approximately 25 ns were not considered in the least square fit as they have some prompt contribution coming from the decay of ^{199}Pt . The resulting value of the Lamor precession frequency ω_L is (40.3 ± 1.2) MHz and the calculated g-factor is $+0.335 \pm 0.010$. In addition to yielding the frequency, the fit also gives information on the amplitude or the effective anisotropy observed which must be satisfactorily accounted for. The theoretical angular correlation coefficient for the cascade, $A_{22}(\text{theo}) = 0.2207$, after appropriate solid angle corrections reduces to a value 0.170 and this is approximately equal to the observed $A_{22}(\text{exp}) = 0.166 \pm 0.015$. The above result is in agreement with the expectations since Pt metal has a cubic structure and no quadrupole interactions should be present to influence the A_{22} . In addition the results show that there is no significant radiation damage caused by the neutron irradiation of the samples.

One can observe in Table I a resemblance in some of the properties of the first $5/2^-$ state in ^{195}Pt , ^{197}Pt , ^{197}Hg and ^{199}Hg . In particular the g-factors of these states are strikingly similar. The enhancement factor for the $5/2^- \rightarrow 1/2^-$ E2 transition is approximately ten or more in all the nuclei and should indicate a predominantly collective character of the state. It was pointed out by de-Shalit⁽¹⁾ that the properties of the first two excited states in ^{199}Hg could be understood in terms of the core excitation model where the $5/2^-$ and $3/2^-$ states are formed by the coupling of the $p_{1/2}$ neutron to the first 2^+ state of the neighbouring even-even nuclei. It was later pointed out by Gal⁽²⁾ that such a simple model could not explain the properties of the states in ^{195}Pt and he suggested some what different model where an admixture of the first two 2_1^+ and 2_2^+ state in ^{194}Pt and ^{196}Pt contribute to the formation of $3/2^-$ and $5/2^-$ doublet. Gal obtained a fairly good agreement in the case of ^{195}Pt for the energies, multipolarities and transition rates. However, with the exception of $5/2^-$ state at 130 keV theory could not explain the observed g-factors of these states. Kalish and Gal⁽⁶⁾ and Vianden and Krien⁽¹¹⁾ have used the similar approach to predict the magnetic moment of the 158 keV and 134 keV $5/2^-$ states in ^{199}Hg and ^{197}Hg respectively with some success. However as pointed out by Gal himself the magnetic moments are extremely sensitive to the admixture of higher configurations and even to small single particle contributions. There are no specific calculations available for the magnetic moment of the 53 keV state in ^{197}Pt , from any of the theories. From the observed resemblance with $5/2^-$ state in ^{195}Pt , ^{197}Hg and ^{199}Hg it may not be too unreasonable to assume a some what similar structure for this state in ^{197}Pt .

Table I

Some Properties of the First $5/2^-$ state in ^{195}Pt , ^{197}Pt , ^{197}Hg and ^{199}Hg

| Energy (keV) | $T_{1/2}$ (ns) | $B(E2)_{\text{exp}}/B(E2)_{\text{s.p.}}$ | g-factor exp | g-factor theo |
|-----------------------|-------------------------|--|-------------------------|------------------|
| ^{195}Pt 130 | $0.87 \pm 0.03^{(4)}$ | $10^{(3)}$ | $0.35 \pm 0.04^{(12)}$ | $0.35^{(7)}$ |
| ^{197}Pt 53 | $16.58 \pm 0.17^{(10)}$ | $10^{(9)}$ | $0.335 \pm 0.010^*$ | |
| ^{197}Hg 134 | $8.066 \pm 0.008^{(8)}$ | $8^{(12)}$ | $0.342 \pm 0.006^{(8)}$ | $0.468^{(11)}$ |
| ^{199}Hg 158 | $2.45 \pm 0.05^{(8)}$ | $18.5^{(2)}$ | $0.352 \pm 0.013^{(8)}$ | $0.480^{(8)}$ |

*Present work

RESUMO

O fator g do estado de 53 keV na ^{197}Pt foi medido utilizando-se o método de Correlação Angular Perturbada Diferencial (TDPAC) gama-gama com um campo magnético externo de 25,1 kG. As medidas foram realizadas usando-se a cascata gama 346-53 keV do decaimento da ^{197}Pt . O valor do fator g obtido foi $+0,335 \pm 0,010$. Este resultado é comparado com fatores g de estados semelhantes da ^{195}Pt e dos $^{197,199}\text{Hg}$.

REFERENCES*

1. DE-SHALIT, A. Core excitation in nondeformed Odd-A nuclei. *Phys. Rev.*, 122:2530-6, 1961.
2. GAL, A. Core excitation in ^{195}Pt . *Phys. Lett.*, 20:414-6, 1966
3. GRODZINS, L.; BORCHER, R. R.; HAGEMANN, G. B. Coulomb excitation in Platinum isotopes. *Nucl. Phys.*, 88:474-80, 1966.
4. HERMATZ, B. Nuclear data sheets. Revised A-chains, A = 195. *Nucl. Data Sheets*, 23:631, 1978.
5. KALISH, R.; BORCHER, R. R.; HUGEL, H. W. Revised level scheme of ^{199}Hg following Coulomb excitation. *Nucl. Phys.*, A161:637-49, 1971.
6. KALISH, R. & GAL, A. The decay of the $13/2^+$ isomeric state and the structure of the low-lying states in ^{199}Hg . *Nucl. Phys.*, A175:652-62, 1971.
7. KISSLINGER, L. S. & SORENSEN, R. A. Spherical nuclei with simple residual forces. *Rev. Mod. Phys.*, 35:853-915, 1963.
8. KRIEN, K.; KROTH, K.; SAITOVITCH, H.; THOMAS, W. Magnetic moments and half lives of the $5/2$ states in ^{199}Hg and ^{197}Hg and the discrepancies in the experimental hyperfine fields at Hg in Fe. *Z. Phys.*, A283:337-343, 1977.
9. MALMSKOG, S. G. The half life of the 53 keV lived in ^{197}Pt . *Ark. Fys.*, 34:195-98, 1967.
10. SOARES, J. C.; MELO, A.; GIL, F. B.; SAXENA, R. N. To be published.
11. VIADEN, R. & KRIEN, K. Quadrupole moment of the $5/2$, 134 keV state in ^{197}Hg . *Nucl. Phys.*, A277:442-50, 1977.
12. WOLBECK, B. & ZIOUTAS, K. The g-factor of the 130 keV state in ^{195}Pt measured by Mossbauer effect. *Nucl. Phys.*, A181:289-97, 1972.

(*) Bibliographic references related to documents belonging to IPEN Library were revised according with NB-66 of Associação Brasileira de Normas Técnicas.

INSTITUTO DE PESQUISAS ENERGÉTICAS E NUCLEARES
Caixa Postal, 11 049 – Pinheiros
CEP 05508
01000 – São Paulo – SP

Telefone: 211-6011
Endereço Telegráfico – IPENUCLEAR
Telex – (011) 23592 - IPEN - BR