

SEAWEEDS AS SOURCE OF THE ESSENTIAL ELEMENTS

C.R. Albuquerque¹. V.A. Maihara². C.B. L. Alves³. P.S.C. Silva⁴

¹Instituto de Pesquisas Energéticas e Nucleares (IPEN / CNEN - SP)
Av. Professor Lineu Prestes 2242
05508-000 São Paulo. SP
calbuuquerque@gmail.com

²Instituto de Pesquisas Energéticas e Nucleares (IPEN / CNEN - SP)
Av. Professor Lineu Prestes 2242
05508-000 São Paulo. SP
vmaihara@ipen.br

³Instituto de Pesquisas Energéticas e Nucleares (IPEN / CNEN - SP)
Av. Professor Lineu Prestes 2242
05508-000 São Paulo. SP
cassiomen@hotmail.com

⁴Instituto de Pesquisas Energéticas e Nucleares (IPEN / CNEN - SP)
Av. Professor Lineu Prestes 2242
05508-000 São Paulo. SP
pscsilva@ipen.br

ABSTRACT

Overtime seaweeds have been used as a food mainly due to their high nutritional value. This type of food is considered as functional food and contributes to the nutritional human requirements, being beneficial to human health. In this study 13 edible seaweed samples acquired in the marked of São Paulo city were analyzed and the concentrations of elements Cl. K. Mg. Mn and Na were determined by Instrumental Neutron Activation Analysis (INAA). The following edible seaweeds were analyzed: Nori (*Porphyra umbilicates*); Hijiki (*Hijikia fusiforme*); Kombu (*Laminaria sp.*) and Wakame (*Undaria pinnatifida*) species from China, USA, Japan and South Korea. The *Undaria pinnatifida* species presented the highest Na concentration and the lowest K level. The highest variation was obtained for Mn in the *Porphyra umbilicates* species.

INTRODUCTION

Marine algae or seaweeds are autotrophic and photosynthetic organisms that differ from plants and may be unicellular or multicellular [1]. It is also the basis of the food chain for many systems, serving as food for zooplankton, which are heterotrophic microorganisms present in plankton. and are staple foods for larger organisms. Thus these algae play a key role in maintaining life on the planet. They are excellent sources of minerals because of the ability to absorb these inorganic substances from the environment and store them in their systems and can be used as biological indicators in many studies [2. 3].

In Asian countries algae have been used for centuries as ingredients in soups and salads as low-calorie foods. while in Western countries their use has traditionally been limited to industrial use as a raw material for extracting alginates from brown algae such as agar- agar

and red algae carrageenan [4]. These polysaccharides have been widely used by the food industry as texture modifiers because of their high viscosity and gelling [5].

Nowadays the use in Western countries is not limited to industrial use, since overtime the influence of oriental culture in the world has increased considerably. especially in cooking. In recent years algae consumption has increased due to the characteristic functional ingredient for formulating healthy foods. aiding in vegan diets to improve nutrition. The production of meat containing seaweed opens new perspectives for the use of these algae [2].

Studies have been carried out to verify the efficacy of algae in alternative treatments of various diseases, such as cardiovascular diseases due to their antioxidant properties [6. 7]. It is known that marine algae contain vitamins, minerals and in particular have high concentrations of long-chain polyunsaturated omega-3s, which reduces the risk of atherosclerosis and inflammatory heart disease [4].

Essential elements are nutrients recognized as fundamental to the functions of the human body and are traditionally divided into macro minerals and micro minerals or trace elements. Macro minerals are required for the human organism in large quantities, since micro minerals or trace elements are needed for the human organism in minor amounts.[8]

In this study four species of edible seaweeds commercialized in São Paulo were analyzed: Hijiki algae (*Hijikia fusiforme*), Kombu (*Laminaria sp*), Nori (*Porphyra umbilicate*) and Wakame (*Undaria pinnatifida*) and the concentrations of the elements Cl, K, Mg, Mn and Na were determined using Instrumental Neutron Activation Analysis (INAA).

This technique is important in food research. providing reliable results for a number of elements that have great nutritional importance [9].

MATERIALS AND METHODS

2.1 Sample Preparation

Thirteen samples from four species of edible seaweeds commercially available to consumers were purchased from super markets all around São Paulo city. The samples were ground and homogenized using a household mixer fitted with titanium blade, until they acquire a powder consistency. Table 1 shows the different species of seaweed and its origins.

Table 1: Identification of seaweeds and their respective origin

Edible seaweed specie	Origin
<i>Porphyra umbilicates</i> (Nori)	China
<i>Porphyra umbilicates</i> (Nori)	Korea
<i>Porphyra umbilicates</i> (Nori)	Japan
<i>Porphyra umbilicates</i> (Nori)	U.S.A.
<i>Hijikia fusiforme</i> (Hijiki)	Japan
<i>Hijikia fusiforme</i> (Hijiki)	China
<i>Hijikia fusiforme</i> (Hijiki)	Korea
<i>Undaria pinnatifida</i> (Wakame)	China
<i>Undaria pinnatifida</i> (Wakame)	Korea
<i>Undaria pinnatifida</i> (Wakame)	China
<i>Laminaria sp</i> (Kombu)	Korea
<i>Laminaria sp</i> (Kombu)	Japan
<i>Laminaria sp</i> (Kombu)	China

2.2. Instrumental Neutron Activation Analysis (INAA)

For the Cl, K, Mg, Mn and Na determination by INAA, about 0.100 g of the seaweed samples were irradiated for 20 seconds in a thermal neutron flux of 1.0×10^{12} n cm⁻² s⁻¹ in a pneumatic station of the nuclear research reactor IEA-R1 at IPEN-CNEN/SP. Primary Cl, K, Mg, Mn and Na standards and biological certified reference material were simultaneously irradiated with the samples for standardization and quality control purposes.

Gamma ray measurements were performed using a GC2018 Canberra HPG e detector coupled to a Canberra DSA-1000 multichannel analyzer. Gamma ray spectra were collected and processed using a Canberra Genie 2000 version 3.1 spectroscopy software. Element content calculations were carried out using a Microsoft Excel spreadsheet.

RESULTS AND DISCUSSION

3.1 Quality control

The analytical quality of the results for the technique used (INAA) was checked for the accuracy and precision of the experimental values obtained for the reference material. Mixed Polish Herb (MPH) from the Institute of Nuclear Chemistry and Technology- (INCT-Poland). The results showed relative errors less than 15% and relative standard deviations below 9%. as can be verified in Table 2.

Table 2: Concentrations of elements in the Certified Reference Material

ELEMENTS	INCT-MPH-2 (Mixed Polish Herb)			
	x ± sd ^a	DPREL%	ER %	Certified Value
Cl %	0.259 ± 0.015	5.7	9.0	0.2840 ± 0.0215
K %	1.747 ± 0.081	4.6	8.5	1.91 ± 0.12
Mg %	0.276 ± 0.041	15	5.7	0.292 ± 0.018
Mn mg/kg	178.9 ± 7.5	4.2	6.3	191 ± 12
Na %	0.0359 ± 0.0019	5.3	-	(0.0350) ^b

^a:media and standard deviation of 2 individual determinations; ^b:informative value

3.2 Results of essential element in edible seaweeds

Thirteen samples belonging to four species of edible seaweeds were analyzed by INAA.

Table 3 shows mean results for the essential elements in seaweeds obtained of duplicate determinations. All results presented are related to the dry weight (dw) of the samples as can be verified in Table 3.

Table 3: Concentrations of the elements in the samples of seaweeds

Species	Origin	µg/kg		%								
		Mn		Cl		K		Mg		Na		Na/K
		Mean ^a	SD	Mean	SD	Mean	SD	Mean	SD	Mean	SD	
<i>Porphyra umbilicates</i> (Nori)	China	26.8	1.6	0.687	0.048	1.445	0.018	0.356	0.019	0.872	0.042	0.61
	Korea	34.5	3	0.518	0.009	1.56	0.14	0.277	0.011	0.685	0.069	0.44
	Japan	13.3	0.7	1.181	0.082	2.5	0.09	0.312	0.024	0.45	0.033	0.18
	USA	33.2	2.3	0.951	0.056	1.86	0.17	0.369	0.017	0.765	0.047	0.41
<i>Hijikia fusiforme</i> (Hijiki)	Japan	21.3	1.6	0.823	0.069	3.53	0.2	0.684	0.064	0.918	0.079	0.26
	China	23.8	2.1	1.88	0.2	1.31	0.14	0.604	0.079	1.29	0.1	0.98
	Korea	20.3	2	0.565	0.055	3.93	0.14	0.49	0.064	1.04	0.14	0.26
<i>Undaria pinnatifida</i> (Wakame)	China	13.9	1.7	Not determined	Not determined	0.481	0.053	0.403	0.04	8.84	0.85	18.4
	Korea	13.2	1.7			0.285	0.07	0.818	0.075	7.03	0.51	24.6
	China	28.7	0.3			0.331	0.007	0.905	0.069	7.30	0.39	22.1
<i>Laminaria sp</i> (Kombu)	Korea	2.4	0.3	9.52	0.45	9.25	0.93	0.536	0.33	2.54	0.18	0.27
	Japan	4.9	0.3	10.6	1.1	8.3	0.48	0.90	0.12	3.37	0.38	0.41
	China	2.2	0.6	6.612	0.056	11.7	1.6	0.325	0.032	2.13	0.19	0.18

The element concentrations in seaweeds are strongly dependent on the types of species, maturity and environmental conditions [4].

In this study large variability can be observed among the seaweed species in relation to their Cl, K, Mg, Mn and Na content. The larger variability was found for Mn in the *Porphyra umbilicates* species from different countries.

However *Hijikia fusiforme*, *Undaria Pinnatifida*, *Laminaria sp* species collected from different countries showed similar levels of the Mg, Mn, Na and K elements.

Chloride content of the seaweed ranged from 0.518 % dry weight in specie *Porphyra umbilicates* (Nori korea) to 1.181 % dw in *Porphyra umbilicates* (Nori-Japan). Houa and Yanb (1998) observed that Cl concentrations in seaweeds are also highly variable and for the same species of algae. However it is not be observed in the analyzed samples.

The *Porphyra umbilicates* samples presented the highest Mn concentrations with range 13.3 to 33.2 mg/kg dw. Mg concentrations in seaweeds not varied over a wide range (0.28 to 0.91 mg/kg dw).

The highest Na concentration was observed in the *Undaria pinnatifida* (7.03- 8.84 %). It is about 10-fold higher than that *Porphyra umbilicates* (0.45 to 0.872% dw) and *Hijikia fusiforme* (0.918 to 1.29 % dw). Similar result was obtained by Dawczynski et al (2007). In contrast *Undaria pinnatifida* showed the lowest K concentrations.

According to Dawczynski et al (2007) the consumption of food with a high Na/K ratio quotient may cause high blood pressure as show in experiment rat. In this study the Na/K ratio are low, about 0.18 – 0.92 to *Porphyra umbilicates*, *Hijikia fusiforme* and *Laminaria sp*. On the other hand the Na/K ratio was high to *Undaria pinnatifida*, circa of the 20-fold than the other analyzed seaweeds (18.4-24.6).

CONCLUSION

INAA proved to be useful to determine the essential elements in seaweeds. The species analyzed showed to be safe due to their low Na/K ratios, except for *Undaria pinnatifida*. For this species the consumption should be moderate.

ACKNOWLEDGMENTS

The authors would like to thank the Institute of Energy and Nuclear Research (IPEN/CNEN) and the CNPq for the fellowship.

REFERENCES

- 1- E. C Vidotti, M. C. E. Rollemberg. “Algas: da Economia nos Ambientes Aquáticos à Bioremediação e à Química Analítica”. *Quimica. Nova*. **27**. 139-145 (2004)
- 2- C. Dawczynski, U. Schäfer, M. Leiterer, And G. Jahreis. ‘Nutritional and Toxicological Importance of Macro, Trace, and Ultra-Trace Elements in Algae Food Products’. *Journal of Agricultural and Food Chemistry*. **55**. 10470-10475 (2007).
- 3- S. R. Rocha, Sanchez-Muniz, F. J. Gómez-Juaristi, M.T.L. Marín. “Trace elements determination in edible seaweeds by an optimized and validated ICP-MS method” *Journal of Food Composition and Analysis*. **22**. 330–336 (2009).

- 4- S.I Cofrades. L. López-López. C Bravo. S. Ruiz-Capillas. M.T Bastida. F. Larrea Juménez-Colmenero “Nutritional and Antioxidant Properties of Different Brown and Red Spanish Edible Seaweeds” *Food Science Technology International*. **16**. 361-370 (2010).
- 5- A.Jiménez-Escrig B.Sc; F.J.Sánchez-Muniz “Dietary fibre from edible seaweeds: Chemical structure, physicochemical properties and effects on cholesterol metabolism” *Nutrition Research* **20**. 585-598. (2000)
- 6- E. Shalaby. “Algae as promising organisms for environment and health” *Plant Signaling & Behavior*. **6**. 1338-1350 (2011).
- 7- C.K. Ku. Y. Yang. Y. Park. J. Lee. “Health Benefits of Blue-Green Algae: Prevention of Cardiovascular Disease and Nonalcoholic Fatty Liver Disease”. *Journal of Medicinal Food*. **16**. 103-111 (2013).
- 8- Gallager. L. Margie.”The nutrients and their metabolism. In: Mahan LK. Escott-Stump S (eds)”. *Krause’s Food nutrition therapy Saint Louis*: Saunders (2008).
- 9- R.P. Avegliano. V.A Maihara. F.F. A Silva. Brazilian ”Total Diet Study: Evaluation of essential elements”. *Journal of Food Composition and Analysis*. **24**. 1000-1016 (2011).
- 10- Xiaolin Houa; Xiaojun Yanb “Study on the concentration and seasonal variation of inorganic elements in 35 species of marine algae” *Science of The Total Environment*. **222**. 141-156 (1998).