

STUDY OF LONG TERM SF₆ MOLE FRACTIONS IN AMAZON AND BRAZILIAN COAST

L.S. Basso¹, R.S. Santos¹, V.F. Borges², L.V. Gatti^{1,2}, L.G. Domingues^{1,2}, C.S.C. Correia^{1,2}, L. Marani¹, S.P. Crispim¹, R.A.L. Neves¹, M. Gloor³, J.B. Miller⁴

¹Earth System Science Center (CCST), National Institute for Space Research (INPE), São José dos Campos, SP, Brazil

²Nuclear and Energy Research Institute (IPEN), SP, Brazil

³School of Geography, University of Leeds, Leeds LS92JT, UK.

⁴Global Monitoring Division, Earth System Research Laboratory, National Oceanic and Atmospheric Administration (NOAA), Boulder, Colorado 80305, USA

Email: luanabasso@gmail.com

1. Introduction

SF₆ is one of the most potent greenhouse gases known, having a very high global warming potential of 23,500 (relative to CO₂). Its surface fluxes include anthropogenic emissions from applications in industry and very minor uptake by the oceans. SF₆ is inert throughout the troposphere and stratosphere and is slowly photolyzed in the mesosphere, resulting in an estimated atmospheric lifetime of 3200 years, so its emissions accumulate in the atmosphere and can be estimated directly from its observed rate of increase^[1]. Its global mole fraction was around 9.5 ppt in 2016, almost twice the level observed in the mid-1990s^[2]. Since it is a very stable gas in the atmosphere, its annual growth rate has been relatively constant since the 1980s, and has been increasing in a very linear way^[3]. Brazil is not an SF₆ producer, therefore, the emissions reported in the Brazilian inventory are due only to leaks in equipment installed in the country due to its maintenance or disposal.

2. Methodology

SF₆ atmospheric measurements were started with vertical profiles using small aircrafts, since 2000 in Santarém (SAN; 2.86°S; 54.95°W), 2010 in Rio Branco (RBA; 9.38°S, 67.62°W), Alta Floresta (ALF; 8.80°S, 56.75°W), Tabatinga (TAB; 5.96°S, 70.06°W) and Tefé (TEF; 3.31°S, 65.8°W, which started in 2013 to replace TAB, results from these sites will be named TAB_TEF), all these sites located in Brazilian Amazon Basin. In 2006, we started flask measurements at Arembepe (ABP, 12,75°S, 38,15°W; between 2006-2009) located at the Brazilian Atlantic coast, and since 2010 started two more locations, Salinópolis (SAL; 0.60°S, 47.37°W) and Natal (NAT; 5.48°S, 35.26°W). The samples from the Brazilian coast were collected weekly by using a pair of glass flasks (2.5L) and a portable sampler. At Amazon sites samples from vertical profiles were collected, generally fortnightly, using a semi-automatic sampling system, which consists of separate compressor and flask units, developed by ESRL/NOAA.

3. Results and Discussion

The SF₆ growth rate obtained for each study site are shown in Table 1, it was observed that in all sites SF₆ mole fractions showed an increase over the studied period, following the global growth rate. SAN results show that SF₆ mole fractions increased since 2000, by nearly 5.3 ppt between 2000 and 2018 (Figure 2), and by a mean increase rate of 0.28 ppt per year.

Figure 3 shows the results obtained during 2010 until 2018 in the four Amazon sampling sites, and the SF₆ global mean mole fractions for this period. It was observed during all these years that mole fractions at all our stations are generally similar to the global mean, with an annual growth ratio between 0.32 and 0.33 ppt/year. Can be observed an annual seasonality, with higher values between January to beginning of May. This seasonality can also be observed in SAL (located in the Brazilian coast). Examination of air parcel paths using HYSPLIT^[4,5] for these periods confirmed that some air parcels arriving at these stations have travelled from the northern hemisphere to the sites. The mean position of Intertropical Convergence Zone (ITCZ) shows variation along the year. Between January and beginning of May the ITCZ is below SAL position, therefore the air masses for SAL and Amazon sites coming from north hemisphere. While in NAT and ABP cannot be observed this seasonality. The air masses back trajectories calculated for NAT and ABP show that the air masses arriving entirely from the South Atlantic Ocean (Figure 3).

4. Conclusions

Results of this long-term measurements showed that all sites had a continuous increase in concentrations, with an annual growth ratio between 0.32 and 0.33 ppt/year (2010-2018), that is similar to the global increase in this period (0.32 ppt/year using the NOAA data). Considering only the SAN measurements (2000-2018) the annual growth ratio is 0.28 ppt/year, the same observed for the global mean concentration during this period.

These results indicate that Amazon and Brazilian northeast coast do not have significant emissions of SF₆ and its mole fractions following the global growth ratio.

Acknowledgments

FAPESP (2016/02018-2, 2008/58120-3, 2011/51841-0, 2018/14006-4), NASA, ERC (GEOCARBON, Horizon 2020/ASICA), NERC (NE/F005806/1), CNPq (480713/2013-8).

References

- [1] Myhre, G., D. Shindell, F.M. Bréon, W. Collins, J. Fuglestedt, J. Huang, D. Koch, J.F. Lamarque, D. Lee, B. Mendoza, T. Nakajima, A. Robock, G. Stephens, T. Takemura, H. Zhang, 2013: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. ISBN: 978-1-107-05799-1.
- [2] WMO, 2018: World Meteorological Organization. *Greenhouse Gas Bulletins. The state of greenhouse gases in the atmosphere using global observations through 2017*. n.14. Disponível em: <
<https://gaw.kishou.go.jp/static/publications/summary/sum42/sum42.pdf>>.
- [3] Kovács, T., W. Feng, A. Totterdill, J.M.C. Plane, S. Dhomse, J.C. Gómez-Martín, G.P. Stiller, F.J. Haenel, C. Smith, P.M. Forster; R.R. García, D.R. Marsh, M.P. Chipperfield, 2017: Determination of the atmospheric lifetime and global warming potential of sulfur hexafluoride using a three-dimensional model. *Atmospheric Chemistry and Physics*, v.17, p.883–898, doi:10.5194/acp-17-883-2017.

- [4] Stein, A.F., R.R. Draxler, G.D. Rolph, B.J.B. Stunder, M.D. Cohen, and F. Ngan, 2015: NOAA's HYSPLIT atmospheric transport and dispersion modeling system. *Bulletin of the American Meteorological Society*, 96, 2059-2077, <http://dx.doi.org/10.1175/BAMS-D-14-00110.1> this link opens in a new window.
- [5] Rolph, G., A. Stein, and B. Stunder, 2017: Real-time Environmental Applications and Display sYstem: READY. *Environmental Modelling & Software*, 95, 210-228, <https://doi.org/10.1016/j.envsoft.2017.06.025>this link opens in a new window. (<http://www.sciencedirect.com/science/article/pii/S1364815217302360>)



Figure 1. Sample sites located in Brazilian Amazon and Brazilian coast

Table 1. SF₆ growth rate to Brazilian Amazon sites and Brazilian coast sites

| Brazilian Coast Sites | | |
|------------------------|---------|------------------------|
| Period | Site | growth rate (ppt/year) |
| 2006-2009 | ABP | 0.27 |
| 2010-2016 | SAL | 0.32 |
| 2010-2018 | NAT | 0.33 |
| Brazilian Amazon Sites | | |
| Period | Site | growth rate (ppt/year) |
| 2000-2018 | SAN | 0.28 |
| 2010-2018 | ALF | 0.33 |
| 2010-2018 | RBA | 0.33 |
| 2010-2018 | SAN | 0.33 |
| 2010-2018 | TAB_TEF | 0.32 |




Figure 2. Temporal series of SF₆ measurements for SAN region (green) and SF₆ global mean mole fractions (black line) from NOAA, between 2000-2018.




Figure 3. Temporal series of SF₆ measurements for Amazon sites (ALF, RBA, SAN, TAB_TEF) and SF₆ global mean mole fractions (black line) from NOAA, between 2010-2018




Figure 4. Temporal series of SF₆ measurements for Brazilian Northeast coast sites (ABP, NAT, SAL) and SF₆ global mean mole fractions (black line) from NOAA, between 2006-2018