

ISSN 0103-6440

ISLD 2004

BRAZILIAN DENTAL JOURNAL

VOLUME 15 (Special issue): SI-01-SI-136, 2004

Dental Foundation of Ribeirão Preto

BRAZILIAN DENTAL JOURNAL

Dental Foundation of Ribeirão Preto

<http://www.forp.usp.br/bdj/bdj.htm>

e-mail: bdj@forp.usp.br

EDITORS

Jesus Djalma Pécora
Paulo Cesar Saquy
Manoel Damião Sousa Neto

ASSISTANT EDITORS

Ricardo Gariba Silva Eduardo Luiz Barbin
Júlio César E. Spanó Melissa Andréia Marchesan

EDITORIAL BOARD

Geraldo Maia Campos	Antonio Luiz B. Pinheiro
Heitor Panzeri	Nicolau Tortamano
Miguel Angel Sala	Maria Fidela de Lima Navarro
Dionísio Vinha	Jaime A. Cury
Jesus Djalma Pécora	Roberto Holland
Delsa Deise Macchetti	Lourenço Correr Sobrinho
Lucy Dalva Lopes	Luis Antônio Salata
Nelo Francisco Romani	Luis César Peres
Regis Alonso Verri	Julio Barrancos Mooney - Argentina
Moacyr da Silva	Julian B. Woelfel - USA
Izabel Ioko Ito	Ivar A. Mjör - Norway
Reinaldo Brito e Dias	Ole Carlsen - Denmark
Ruberval A. Lopes	Lief Tronstad - Norway
Esther Goldenberg Birman	Walter A. Bretz - USA
Flávio Fava de Moraes	Benjamin Martinez - Chile
Aldo Brugnera Júnior	A.D. Walmsley - Birmingham, UK
Vera Cavalcanti de Araújo	Harvey Wigdor - USA
Arthur Belem Novaes Jr.	Benjamin Peretz - Israel
Carlos Estrela	Trevor F. Walsh - UK
Léa Assed Bezerra da Silva	Harald J. Schiel - Switzerland

DENTAL FOUNDATION OF RIBEIRÃO PRETO

DIRECTORS

Jesus Djalma Pécora
Ricardo Gariba Silva
Iara Augusta Orsi

The Journal is indexed by Medline, PubMed, SciELO, DEDALUS and ERL.

Financial assistance for the publication of the Journal provided by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and FAPESP.

Support to Scientific Publications Program

MCT  CNPq  FINEP FAPESP

Poster Presentation

296.67 m to 200 and 300 mJ, respectively). The volume also increased when 300 mJ was used. Our finding indicated that cavities obtained were round or slight oval. The diameter was not affected by energy level, however high energy level increased depth and volume of the cavities.

P39

Comparison between photodynamic therapy and a bactericidal solution in the treatment of dental alveolitis microbiological evaluation.

Hayek, R.R.A.; Yamada Júnior, A.M.; Garcez, A.S.; Nuñez, S.C.; Suzuki, L.C.; Ribeiro, M.S..
Mestre em Ciências, IPEN/USP.

Photodynamic therapy is a new therapeutic modality to treat microbial infections. The aim of this pilot study was to compare this new therapeutic approach with chlorhexidine in mini pigs induced alveolitis. Alveolitis is an inflammatory disease that involves the superficial bone layer in the dental alveolus. Eight dental elements were extracted from mini pigs and immediately after the alveolus were contaminated with *Streptococcus aureus*. After seven days the area presented edema and hyperemia. In the chlorhexidine group the alveolus were debrided and irrigated with a 0.12% solution of chlorhexidine. In Laser group the alveolus were debrided and photodynamic therapy was performed consisting in the application of a azulene paste inside the alveolus with a pre-irradiation time of 5 minutes followed by irradiation with a low power diode laser, $\lambda = 685$ nm, P= 50 mW, E= 9 J, for 3 minutes. Microbiological samples were harvested before and after treatment for both groups with sterile paper points. The results showed bacterial reduction in both groups, although photodynamic therapy was significantly more effective to reduce the population of *S. aureus*. This finding indicates that photodynamic therapy can be an alternative method to the treatment of alveolitis.

P40

Effects of the diode laser irradiation on root surfaces. thermal analysis.

Haypek, P.; Theodoro, L.H.; Bachmann, L.; Eduardo, C. P.; Sampaio, J.E.C.; **Zezell, D.M..**
Doutoranda em Dentística Restauradora- Faculdade de Odontologia da Universidade de São Paulo-USP.
The purpose of this study was to evaluate the temperature rising

inside the pulpar cavity during the diode laser irradiation on the root surface. Twelve single root teeth were used and their mesial and distal surfaces were irradiated (808 nm, 400 m, 30 s), in a continuous mode (CW) and in an switched mode (SW) (10 Hz). The temperature rise was evaluated depending with the power used (0.4 W to 2.2 W). The temperature was monitored with a termopar (T) fixed inside de pulpar cavity during the irradiation. In a second step two parameters was used in a SW (10 Hz). The safety temperature was keep using the power: Group A - 0.9 W (mesial) and Group B - 1.08 W (distal). The results showed the termal pulpar events (T) during the irradiation, based in an specific relation with the power (P), like $T = -0.4 + 3.7P$ in a CW and $T = -0.2 + 2P$ in a SW. The temperature rising during the irradiation was $3.4 \pm 0.4^\circ\text{C}$ on Group A and $4.0 \pm 1.0^\circ\text{C}$ on Group B. The irradiation mode and the power are much important to temperature rise inside the pulpar cavity and the 0.9 W and 1.08 W radiation on the root surfaces were safety.

P41

Effects of Nd:YAG laser irradiation to fauces.

Hotta, K.; Yamaguchi, H.; Sekine, A.; Kobayashi, K.; Gomi, K.; Arai, T..

Department of Periodontics and Endodontics Tsurumi University School of Dental Medicine. Yokohama , Kanagawa, Japan Aims:

The aim of this study was to evaluate in vivo systemic blood pressure and pulse rate during Nd:YAG laser irradiation of fauces (Mandibular fauces was distributed in glossopharyngeal nerve of Rr.linuales :parasympathetic nerve). Material and Methods: The 6 adult volunteers participated in this study. The experimental site for each patient was mandibular fauces. Mandibular fauces area was irradiated with Nd: YAG laser irradiation (30 sec at energy level of 100mJ pulses at 15pulses/sec). The systemic blood pressure and pulse rate were monitored for each patient's throughout the laser irradiation procedure. Results: No significant change in systemic blood pressure and pulse rate were found in laser irradiation of fauces. Discussion and Conclusion: Systemic blood pressure and pulse rate were affected for the irradiation to