

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/374256847>

Examination of Natural Radioactivity Concentration and Radiological Exposure of Soil Within Mining Site in Umuahia South Abia State Nigeria, Using High Purity Germanium (HPGe) Gamm...

Article · January 2023

DOI: 10.18576/jrma/080304

CITATIONS

0

READS

67

9 authors, including:



Moses Onudibia

Federal University Wukari Taraba/Instituto de Pesquisas Energéticas e Nucleares (IP...

1 PUBLICATION 0 CITATIONS

SEE PROFILE



Paulo Sergio Cardoso Silva

University of São Paulo

78 PUBLICATIONS 674 CITATIONS

SEE PROFILE



Guilherme Zahn

Instituto de Pesquisas Energéticas e Nucleares

82 PUBLICATIONS 189 CITATIONS

SEE PROFILE



Alloysius Chibuike Ogoto

Federal University Wukari

84 PUBLICATIONS 445 CITATIONS

SEE PROFILE

Examination of Natural Radioactivity Concentration and Radiological Exposure of Soil Within Mining Site in Umuahia South Abia State Nigeria, Using High Purity Germanium (HPGe) Gamma Ray Spectrometry

Moses Ejike Onudibia^{1,2*}, Paulo Sergio Cardoso da Silva², Aniesua Akpan Essiett³, Guiherme Soares Zahn², Frederoco Anthonio Genezini², Edet Essien Imeh³, Ngene Chibuike Nnamani⁵, Alloysius Chibuike Ogodo⁶, Celestial Bede Mfom⁴ and Franca Oyiwoja Okoh¹

¹Department of Pure and Applied Physics, Federal University Wukari, Taraba State P.M.B 1020 Kasinala Road, Wukari, Taraba State, Nigeria.

²Institute for Energy and Nuclear Research (IPEN), Research Reactor Center (CRPq), University of Sao Paulo (USP), São Paulo, Brazil.

³Department of Physics, University of Uyo, Uyo, Akwa-Ibom, Nigeria.

⁴Department of Physics, Federal University of Technology, Ikot Abasi, Akwa Ibom State, Nigeria.

⁵Department of Statistics, Ahmadu Bello University, Zaria, Kaduna, Nigeria.

⁶Department of Microbiology, Federal University Wukari, Taraba State P.M.B 1020 Kasinala Road, Wukari, Taraba State, Nigeria.

Received: 02 Jul 2023, Revised: 04 Jul 2023, Accepted: 24 Aug. 2023.

Published online: 1 Sep 2023.

Abstract: Mining activity is one of the major sources of exposure to radiation. The main goal of this study was to determine the natural radioactivity level and its radiological exposure risk in mining site Umuahia South Abia State, Nigeria, using High Purity Germanium (HPGe) gamma ray spectrometry. The activity concentration of ^{40}K , ^{226}Ra and ^{232}Th in the soil ranges from 31 ± 2 to $367 \pm 27 \text{ Bq kg}^{-1}$, with mean value of $142 \pm 11 \text{ Bq kg}^{-1}$; from 26 ± 2 to $65 \pm 5 \text{ Bq kg}^{-1}$, with mean value of 49 ± 4 ; and from 53 ± 9 to $109 \pm 17 \text{ Bq kg}^{-1}$, with mean value of $77 \pm 16 \text{ Bq kg}^{-1}$, respectively. The activity concentrations of ^{226}Ra and ^{232}Th are above the global values of 32 and 45 Bq kg^{-1} while ^{40}K is depleted in relation to the global mean of 412 Bq kg^{-1} . Radiological parameters are generally in agreement with the values observed for other Nigerian and other countries soil reported in literature. The main responsible for the values observed for the radiological parameters is ^{232}Th followed by ^{226}Ra . The use of this soil for dwelling construction is unlikely to pose any radiological harm for the residents.

Keywords: Natural Radioactivity, Radiological Parameters, Exposure Indices, Soil, Mining, High Purity Germanium (HPGe) Gamma Ray Spectrometry.

1 Introduction

Soils are the actual store house of radionuclide naturally occurring on the Earth's crust responsible for transferring these elements into biological systems [1]. Radioactivity is an observable fact that is connected with energetic atomic nuclei that are automatically decomposed releasing beta, alpha, and neutron particles, or electromagnetic radiation in the form of gamma rays [2]. Soil radioactivity is gaining great scholar attention widely as it one of the main factors in public doses and may help predict changes in the

environment [3,4]. Soil characteristics, geological formations and human practices are important factors for potentially elevating the concentrations of background natural radiation [5]. The level of natural radioactivity in an area depends on the soil types, rocks types and its geology [4]. Moreover, the distribution and variation of radionuclides in surface soil relies on the composition and distribution of radioelements in the bedrock, their physical properties and mechanical properties such as porosity, permeability [5]. Soil, rocks and minerals commonly

*Corresponding author e-mail: mosesmarke@gmail.com

contains naturally occurring radioactive materials (NORM), with half-lives as long as the Earth's age, known as primordial radionuclides [2]. The main radio nuclides of this category are ^{40}K , and the generators of decay series ^{232}Th and ^{238}U .

The radioactivity in the environment needs attention because of the potential harmful effects of the ionizing radiations on tissues of living organisms, though their occurrence is uncommon. Hence, scholars examined natural environmental radioactivity in soil to carry out background study and detect environmental radioactivity levels [6].

Urbanization, mining activities, waste disposal systems, and several human activities are contributors to the radioactivity of the environment [3, 4], when soil, rock and dust material that contains these NORM are spread from the mine areas, during mining and processing, to the surrounding area where they can be amassed in the food chain [6].

Of interest for this study, mining activities and the processing of natural resources have impacted strongly on human being and the environment [5] because NORM is largely responsible for the external or internal irradiation dose received by workers and neighbor population of the mining sites [6]. Internal exposure takes place via the intake of terrestrial radionuclides through inhalation or ingestion [7-8].

In Nigeria, mining activities for coal, lignite, lithium, bitumen, iron ore, and clay, among others, go on mostly by a domestic and underdeveloped mining industry. In the country there are about 44 different types of minerals that have been identified in over 500 locations [9]. Among the minerals, clay holds significant contributions due to its high demand for construction, cement building, and block industry [10]. In Umuahia South, Abia State, South East, Nigeria, present study area, where clay mining goes on, people are involved in mining activities as their source of income and subjected to the unavoidable NORM emission which aids the constant release of radionuclides in the environment. The main goal of this paper is to study the natural radioactivity in the soil on the Umuahia mining sites and to ascertain radiological parameter and the hazard indices, therefore comparing with international organisation recommendations and global average.

2 Materials and Methods

2.1 Study Area

The study area is Umuahia South, South East, Nigeria, where clay mining is going on. Umuahia South is one of the local government areas of Abia State and it has a geographic coordinate of $5^{\circ} 31' 0''$ North, $7^{\circ} 26' 0''$ mean area of 140 km^2 and population of 138,570 [11]. The mining site is about 10 km from Umuahia main town and about 45 km to Aba, Enyimba City of Abia state. The average annual temperature is 26.4°C . About 2333 mm of

precipitation falls annually. The vegetation is ordinarily considered part of tropical rainforest which is the dominant natural vegetation in most parts of southern Nigeria [12].

2.2 Sampling

All of the 15 soil samples were collected using a soil auger sampler at a depth of 5-15 cm, for every sampling point about 1 kg to 1.5 kg were collected. The soil samples were lowered into labelled plastic bags and every bag was properly tied [13].

2.3 Samples Preparation

After the collection of samples from the field, they were kept in natural air-dried in a room at ambient temperature for about one month. Moreover, at the laboratory, the samples were air dried in trays for 7 days and then oven dried at a temperature of 105°C for between 3-4 hours until the samples were well dried with a constant weight [14], at the central lab of Federal University Wukari, for preparation. Soil samples were pounded using mortar and pestle. A sieve of 2mm pore size mesh was used to obtain uniform particle sizes, as recommended by IAEA [15]. The sieved prepared soil samples were fled to Nuclear and Energy Research Institute (IPEN) or Instituto de Pesquisas Energéticas e Nucleares (IPEN), Sao Paulo Brazil for analysis.




Fig.1: Location of the studied area. A) Nigeria; B) Umuahia region; C) Mine sampling points.

2.4 Analysis of Samples

In this work, the radionuclides ^{40}K , ^{228}Ra , and ^{232}Th were determined using p-type High Purity Germanium Detectors (HPGe), Canberra (USA) with relative efficiency $\geq 20\%$, and resolution $\leq 2.00 \text{ keV}$ at 1.33MeV . The activity concentration of the radionuclides was determined using

the following energies: ⁴⁰K was determined using the 1460 keV gamma-ray peak; ²²⁶Ra was determined using the 351 keV (²¹⁴Pb) and 609 keV (²¹⁴Bi) energies, and ²³²Th was determined using 727 keV (²¹²Pb) and 238 keV (²¹²Bi). The samples were counted for 86,400 seconds, background was counted for 172,800 seconds while the certified reference materials (CRM) IAEA-RGU-1 uranium ore, IAEA-RGTh-1, thorium ore and IAEA-RGK-1, potassium sulfate, were counted for 7,200 seconds. Activity concentrations were determined via a punctual calibration comparing the intensity of peak in samples to the respective peaks of the standard reference material. For certainty and to ensure the quality assurance of this work, the IAEA-327 reference material (Radionuclides in Soil) from IAEA was used [16]. The obtained values were 626 ± 43 Bq kg⁻¹ for ⁴⁰K, 36 ± 3 Bq kg⁻¹ for ²²⁶Ra and, 38 ± 7 Bq kg⁻¹ for ²³²Th. All the obtained result agrees with the 95% confidence interval of the reference material: 612 – 630, 32.7 – 35.5 and 37.2 – 39.2 Bq kg⁻¹, for ⁴⁰K, ²²⁶Ra and ²³²Th, respectively.

2.5 Activity Concentration

The activity concentration of the radionuclides has been calculated using the expression Equation 1 [17].

$$A = \frac{N e^{\lambda t_0}}{\epsilon F_{\gamma} M t_c} \quad (1)$$

Where, A is the activity concentration of the radionuclide; N is number of counts at a given energy, subtracted from the BG; λ is the decay constant; t_c is the time of counting; t₀ is the time difference between sampling and start of count; M is the sample mass; ε is the efficiency; F_γ is the emission probability and F_c is the correction factor.

2.6 Radiological Parameters (RP) and Exposure Indices (EIs)

Radiological parameters of exposure indices are commonly used to assess the radiation exposure originating from the environment and building material [1, 18, 19, 20]. The indices calculated in this work are listed and defined below, denoting with Ra_{At}, Th_{At} and K_{At} the activity concentration of ²²⁶Ra, ²³²Th and ⁴⁰K respectively, obtained as described in Equation 2 to Equation 7.

Radium Equivalent Activity: $Ra_{eq} = 0.077 K_{At} + Ra_{At} + 1.43 Th_{At}$ (2)

Absorbed Dose Rate in Air: $AD = 0.0417 K_{At} + 0.462 Ra_{At} + 0.604 Th_{At}$ (3)

Representative Level Index: $RLI = \frac{K_{At}}{1500} + \frac{Ra_{At}}{150} + \frac{Th_{At}}{100}$ (4)

Annual Effective Dose Equivalent: $AEDE = D \times OF_t \times CF_t$ (5)

$$AEDE_{OUT} = D \times 0.7 \times 0.2 \times 8760 \times 10^{-6} \quad (5a)$$

$$AEDE_{OUT} = D \times 0.7 \times 0.8 \times 8760 \times 10^{-6} \quad (5b)$$

Annual Gonadal Equivalent Dose: $AGDE = 0.314 K_{At} + 3.09 Ra_{At} + 4.18 Th_{At}$ (6)

Excess Lifetime Cancer Risk: $ELCR = AEDE \times MLE \times RF$ (7)

Radium Equivalent Activity (Ra_{eq}) represents a weighted sum of the activities of ²²⁶Ra, ²³²Th, and ⁴⁰K in the building materials samples, and is defined as the activity concentration of a radionuclide equivalent to 370 Bq kg⁻¹ of ²²⁶Ra that gives outdoors an external effective dose rate of 1 mSv y⁻¹. So, the Ra_{eq} index should not exceed 370 Bq kg⁻¹ for building materials [21, 22]. The Absorbed Dose Rate in Air (AD) is employed in measuring the exposure that enables ascertaining the quantity of radiation received by the human body due to the concentrations of radionuclides (²²⁶Ra, ²³²Th, and ⁴⁰K) in soil [21].

The Representative Level Index (RLI) expresses the extent the soil materials from a site of interest is convenient for using as construction material [23]. It is also applied to estimate the rate of γ-radiation risks in connection with natural radionuclide in investigated samples (3, 6).

The Annual Effective Dose Equivalent (AEDE) computes the annual effective dose for human beings exposed of a certain dose rate [21, 22]. The AEDE can be obtained by converting the total absorbed dose and the product of occupancy factor (0.2 for outdoor and 0.8 for indoor exposure), of a year, expressed in seconds.

Annual Gonadal Equivalent Dose (AGDE) is used to evaluate the potential effects of exposure to radiation on certain important organs, such as gonads, the activity bone marrow and the bone surface cells since those are the most sensitive parts of the human body [23-24].

Excess Lifetime Cancer Risk (ELCR) is an index related to the risk of developing cancer as a result of amassed exposure to a toxic substance liable to the existence time of human being for a period of lifetime [19].

3 Results and Discussion

3.1 Activity Concentration of Soil the Mining Site

In Tables 1 and 2 the activity concentration of ⁴⁰K, ²²⁶Ra and ²³²Th determined in the soil samples of the present study and comparison of activity concentration of radionuclides of soil (Bq kg⁻¹) determined in the literature, respectively, are presented. The activity concentration of ⁴⁰K, ²²⁶Ra and ²³²Th in this study ranges from 31 ± 2 to 367 ± 27 Bq kg⁻¹ with mean value of 142 ± 11 Bq kg⁻¹; 26 ± 2 to 65 ± 5 Bq kg⁻¹ with mean value of 49 ± 4 and 53 ± 9 to 109 ± 17 Bq kg⁻¹ with mean value of 77 ± 16 Bq kg⁻¹,

respectively.

The activity concentrations ratio $^{226}\text{Ra}/^{40}\text{K}$, $^{232}\text{Th}/^{40}\text{K}$ and $^{232}\text{Th}/^{226}\text{Ra}$ are shown in Figure 2. The point 16, in the figure, were calculated using the mean values for these radionuclides in soil, according to [25]. It can be seen that ^{40}K is depleted in relation to the reference, while ^{232}Th and ^{226}Ra ratio present the same ratio value (around 1.5) of the global mean. This can indicate that ^{232}Th and ^{226}Ra have the same distribution patten in these soil samples and that the ^{226}Ra activity concentration must be approximately equal to that of ^{238}U considering the crustal Th/U activity ratio with an average of about 1.1 [26]. The ^{40}K depletion must be related to the soil type and the predominant weathering occurring in the region forming the soil known as ferruginous tropical [27, 28].




Fig. 2: Activity ratios of the determined radionuclides in soil samples. Point 16 was calculated using the UNSCEAR [25] global average values.

Figure 3 shows the correlation coefficients between ^{40}K , ^{226}Ra and ^{232}Th . It can be seen that ^{40}K do not show significant correlation with ^{226}Ra and ^{232}Th , indicating they probably are linked to different mineral phases or granulometric fractions. Confirming the results showed in Figure 2, the correlation coefficient (Figure 3) between ^{226}Ra and ^{232}Th indicates a strong correlation ($r = 0.74$).

The strong positive correlation between ^{226}Ra and ^{232}Th in the samples indicates that their rock content must be mostly influenced and controlled by the similar source origin [29], and the fact that radium (from the ^{238}U -series) and thorium decay series occur combined together in nature [30].




Fig.3: Linear regression and correlation coefficient ^{40}K x ^{226}Ra , ^{40}K x ^{232}Th , and ^{226}Ra x ^{232}Th

The observed values of ^{226}Ra and ^{232}Th in these samples were higher than global average while the ^{40}K activity concentration had lower value (3). The increasing order of the activity concentrations for the analysed radionuclides is $^{226}\text{Ra} < ^{232}\text{Th} < ^{40}\text{K}$. Comparing the present study with local and multinational related reports, as depicted in Table 2, the mean activity concentration of ^{40}K of similar research is higher than the present study except that of [31] also located in the South-East Enugu region of Nigeria. The ^{226}Ra activity concentrations measured in Nigerian research reports indicate a variation ranging from 12 to 66 Bq kg^{-1} , which is in agreement with the values here reported. The same is also observed in the comparing with international research studies whose variation ranges from 22 to 86 Bq kg^{-1} .

3.2 Radiological Parameters (RPs) and Exposure Indices (EIs)

Table 3 depicts the RPs and EIs. The AD values ranged from 70 ± 8 to 105 ± 12 (nGyh^{-1}), with almost all the samples presenting higher values than the global average with mean values ranging from 50 to 59 nGy h^{-1} [25]. According to the same previous reference, 90% of the world population are exposed to about 20 to over 100 nGy h^{-1} , indicating that the analyzed soil is in well agreement with the absorbed dose registered in most of the world. The annual outdoor and indoor AEDE ranged from 0.06 ± 0.01 to 0.13 ± 0.01 mSv y^{-1} and 0.23 ± 0.03 to 0.52 ± 0.06 mSv y^{-1} , respectively, and are in good agreement with the global average of 0.08 and 0.42 mSv y^{-1} . Radium equivalent presented a variation from 106 to 236 Bq kg^{-1} and all the samples presented values lower than the recommended value of 370 Bq kg^{-1} indicating that this soil used as construction material will not pose unnecessary exposures for the population that can result in late deterministic effects [32]. The indoor ELCR for the soil samples in this study ranged from 0.17×10^{-3} to 0.39×10^{-3} which is very close to the world average of 0.29×10^{-3} . The RLI in the soils analyzed in this study varied from 0.74 to 1.67 and is above the reference value of 0.46 mSv y^{-1} [33].

The model used for EGDE calculation, for a given material be used as construction material, consider a typical cavity type house with infinitely thick walls, which makes it possible to compare the AGED of a house containing concentration of ^{226}Ra , ^{232}Th , and ^{40}K equal with the world average in soil with those obtained using that given material [34]. The value of AGDE considering the global mean of the activity concentrations of ^{226}Ra , ^{232}Th , and ^{40}K is 416 $\mu\text{Sv y}^{-1}$. The calculated AGDE for samples from 320 to 732 $\mu\text{Sv y}^{-1}$, slightly higher than the recommend value

[35] Principal component and classification analysis was applied to the results of radionuclides determined activity concentrations and calculated radiological parameters. The result is shown in Figure 4 where the projection of the cases as a function of factors 1 and 2 are presented. The samples in the right side, with positive loading factor, in the factor 1, are those presenting the lowest ^{226}Ra and ^{232}Th activity concentrations and were collected in places near the border of the mine. Samples located in the left side of the Figure 4, have negative loading factors and present the higher activity concentration for ^{226}Ra and ^{232}Th .

The projection of the variables, determined activity concentrations and calculated radiological parameters, as a function of factors 1 and 2 are presented in Figure 5 showing that ^{40}K is negatively correlated with ^{226}Ra and ^{232}Th and that all the calculated radiological indices are dependent of the activity concentration of the last two nuclides.




Fig. 4: Factors obtained in component analysis showing the loading factor for Factor 1 x Factor 2




Fig. 5: Correlation between activity concentrations ^{226}Ra , ^{232}Th , and ^{40}K and the radiological indices

Table 4 presents radiological indices values of some studies from Nigeria and other world locations. For AD the present study was higher than reported values except in Tanzania and in Nigeria (Anka). The RLI value of the present study was higher than those reported values, except for the value in Al-Khor, Qatar. ELCR value found for the samples of this study is in the same order of others. AGED in present study was lower than the value registered in Tamilnadu, India, and higher than the value reported for Ibadan Oya, in Nigeria. Hence, the mining activities in the local government may influence the radiological characteristics of the Umuahia region soil and it is recommended that constant monitoring of the mining activities and workers be adopted. If used as construction material for brick production, it is unlikely that any radiological harm may be posed by these samples.

Table 1. Activity of soil Umuahia, SUB (BqKg^{-1}).

SID	^{40}K	^{226}Ra	^{232}Th
SUB1	109.72±8.48	43.31±3.35	66.25±8.62
SUB2	366.59±27.03	65.01±4.89	99.49±13.71
SUB3	BDL	54.73±4.47	83.37±10.93
SUB4	111.61±9.49	47.93±4.32	84.99±11.77
SUB5	55.81±4.47	26.06±2.04	53.05±8.62
SUB6	133.82±10.17	39.25±2.95	61.65±8.61
SUB7	47.93±5.02	61.26±5.38	76.15±10.61
SUB8	96.4±8.16	54.27±4.37	109.43±16.89
SUB9	49.21±4.89	56.78±5.06	87.48±87.48
SUB10	326.37±25.58	43.56±3.28	82.64±11.95
SUB11	279.04±22.13	59.97±5.21	78.33±10.33
SUB12	31.45±2.34	38.4±3.08	57.09±7.45

SUB13	119.92±9.18	57.69±4.36	87.63±15.47
SUB14	48.35±5.04	45.28±4.08	60.23±8.93
SUB15	218.71±15.53	40.31±3.54	72.18±11.09
MEAN	142.49±11.25	48.92±4.02	77.33±16.16
GLOBAL MEAN	412	32	45

Table 2. Comparison of soil (BqKg^{-1}) with other scholar's work.

LOCATIONS	^{40}K	^{226}Ra	^{232}Th	REFERENC E
Nigeria (Abia, Umuahia South)	142.49±11.25	48.92±4.02	77.33±16.16	Present Study
Brazil (State of Parana) Soil 1	237 ± 15	50 ± 9	48 ± 3	[36]
Brazil (State of Paraná) Soil 2	258 ± 17	23 ± 4	23 ± 1	
India (Punjab and Himachal Pradesh)	143.04	56.74	87.42	[37]
Iraq	335.8	21.7	9.4	[19]
Turkey	771.57	85.75	51.08	[38]
Nigeria (Itaganmodi, south-western)	505.5	55.3	26.4	[4]
Nigeria (Zamfara)	426.51	12.12	60.117	[39]
Nigeria (Kaduna)	459.56	62.28	155.36	[20]
Nigeria (Niger Delta)	210	18	22	[40]
Nigeria (Alade, Southwestern)	383.77	20.82	16.32	[41]
Nigeria (Ijero Town)	635.41	42.02	43.27	[23]
Nigeria, South-East (Enugu)	100.65	33.16	77.68	[31]
Nigeria, North West (Anka)	380.34	41.60	151.15	[42]

Table 3. Radiological parameters for soil Umuahia.

Sample	AD (nGyh^{-1})	Raeq (BqKg^{-1})	RLI	AEDE _{out} (mSv y^{-1})	AEDE _{in} (mSv y^{-1})	ELCR $\times 10^{-3}$ (mSv y^{-1})	AGDE ($\mu\text{Sv y}^{-1}$)
SUB1	65±7	146±16	1.02±0.11	0.08±0.01	0.32±0.03	0.24±0.03	445±49
SUB2	105±12	236±27	1.67±0.19	0.13±0.01	0.52±0.06	0.39±0.04	732±81
SUB3	76±9	174±20	1.2±0.14	0.09±0.01	0.37±0.04	0.28±0.03	518±59
SUB4	78±9	178±22	1.24±0.15	0.1±0.01	0.38±0.05	0.29±0.04	538±66
SUB5	46±6	106±15	0.74±0.1	0.06±0.01	0.23±0.03	0.17±0.02	320±44
SUB6	61±7	138±16	0.97±0.11	0.07±0.01	0.3±0.03	0.23±0.03	421±48
SUB7	76±9	174±21	1.2±0.15	0.09±0.01	0.37±0.04	0.29±0.03	523±62
SUB8	95±13	218±29	1.52±0.2	0.12±0.02	0.47±0.06	0.36±0.05	655±87
SUB9	81±55	186±130	1.29±0.91	0.1±0.07	0.4±0.27	0.30±0.21	557±383
SUB10	83±10	187±22	1.33±0.16	0.1±0.01	0.41±0.05	0.31±0.04	583±68
SUB11	86±9	193±22	1.37±0.15	0.11±0.01	0.42±0.05	0.32±0.04	600±66
SUB12	54±6	122±14	0.85±0.1	0.07±0.01	0.26±0.03	0.20±0.02	367±41
SUB13	85±12	192±27	1.34±0.19	0.1±0.01	0.41±0.06	0.32±0.04	655±81
SUB14	59±7	135±17	0.94±0.12	0.07±0.01	0.29±0.04	0.22±0.03	556±51
SUB15	71±9	160±21	1.14±0.14	0.09±0.01	0.35±0.04	0.27±0.03	583±62
Mean	75±12	137±28	1.19±0.2	0.09±0.01	0.37±0.06	0.28±0.05	530±83

Global Mean	59	370	1	0.080	0.42	0.29	416
-------------	----	-----	---	-------	------	------	-----

Table 4. Comparison of radiological parameter of soils of present study and other works.

Location	AD (nG yh ⁻¹)	Ra _{eq} (Bq kg ⁻¹)	RLI (Bq kg ⁻¹)	AEDE (mSv y ⁻¹)	ELCR	AGED	Reference
Present work	75±12	137±28	1.19±0.2	0.09±0.01	0.23±0.05	417±83	Present Study
Egypt (Suez Canal)	18.88	39.24	0.30	0.020			[43]
Iraq	23.27	46.82	0.359	0.142			[19]
India (Tamilnadu)	60.73	133.14	0.96	0.075	0.261	744.7	[44]
Tanzania	106.8	230.6	0.831	1.07			[6]
Iraq (Karbala)	29.7	61.11	0.463	0.03	0.12		[19]
Turkey	45	96		0.056			[45]
Qatar (Al-Khor)	22	46		0.026	0.941		[46]
Nigeria (Ibadan Oya)	42.0	90.97	0.62	0.36		284.94	[2]
Nigeria (Anka)	121.8	273.1		0.149			[42]

5 Conclusions

The NORM radionuclides ⁴⁰K, ²²⁶Ra and ²³²Th, resulting from clay mining activities, in the soil mining site in Umuahia South Abia State Nigeria were determined. Compared to the global average, the activity concentrations of ²²⁶Ra and ²³²Th are above and ⁴⁰K are below the mean values. Potassium is depleted in this soil samples, probably as a result of the bad rock weathering that occurs in the region. The activity ratio ²³²Th/²²⁶Ra indicates that these radioactive elements must have the same provenance in the soil formation and link to the same mineral phases. On the other hand, ⁴⁰K did not present correlation with ²²⁶Ra and ²³²Th. Comparing with published values for other Nigerian sites and other countries, ²²⁶Ra and ²³²Th was found in agreement with the range of values reported in literature and the low values measured for ⁴⁰K is also observed in reports for the same region, i.e, Umuahia South, South East, Nigeria. Samples with the highest activity concentrations tend to be concentrated more in the centre of the mining area.

The radiological parameters calculated indicated values slightly above global means for absorbed dose and annual gonadal equivalent dose with ²³²Th responding for the higher contribution, followed by ²²⁶Ra. It was observed that ⁴⁰K do not present correlation with the radiological indices. The results indicate that use of this soil samples for brick production to be used as construction material is unlikely to cause any radiological harm for the individuals.

Acknowledgement

The authors mmensely appreciate the staff of Instituto de Pesquisas Energéticas e Nucleares (IPEN), Centro do reator de pesquisa (CRPq), USP, Brazil and Tertiary Education Trust Fund (TETFUND for their support.

References

- [1] L. Garba, I.O.B. Ewa, M.O.A., Oladipo, and C.U. Omeje, Rare earth element distribution patterns in the soils around the Nigerian Research Reactor-1, [nirr-1] in Zaria, Nigeria, *European Scientific Journal*, **8(16)**, 245-251, 2012.
- [2] D.A. Ayeni, and F.M. Adebisi, Evaluation of natural radioactivity and radiation hazards of soils around petroleum products marketing company using gamma ray spectrometry. *Tanzania Journal of Science*, **48(2)**, 304-312, 2022.
- [3] United Nations Scientific Committee on the Effects of Atomic Radiation, Sources and Effects of Ionizing Radiation, UNSCEAR 2000 Report Vol.1. to the General Assembly, with scientific annexes, United Nations Sales Publication, United Nations, New York., **1**, 2000.
- [4] A.K. Ademola, Radiological risks from natural radionuclides in surface soil of Agbara industrial area, Ogun State, Nigeria, *J. Appl. Sci. Environ Manage*, **25(5)**, 733-739, 2021.
- [5] S. Alashrah, A. El-Taher, H. Mansour., Assessment of radiological parameters and metal contents in soil and stone samples from Harrat Al Madinah, Saudi Arabia. *MethodsX* **5**, 485-494. 2018
- [6] M.S., Gurisha, A.A., Makoba, F.P., Banzi, and H.E.

- Kiyengo, Determination of natural radioactivity levels and radiation hazards from the soil of tin, mining in Kyerwa District, Tanzania. Research Square, 1-15, 2022.
- [7] HA Awad, HMH Zakaly, AV Nastavkin, AM El Tohamy, A. El-Taher., Radioactive mineralizations on granitic rocks and silica veins on shear zone of El-Missikat area, Central Eastern Desert, Egypt. *Applied Radiation and Isotopes* 168, 109493. 2021
- [8] AE Abdel Gawad, MM Ghoneim, A El-Taher, AA Ramadan ., Mineral chemistry aspects of U-, Th-, REE-, Cu-bearing minerals at El-Regeita shear zone, South Central Sinai, Egypt. *Arabian Journal of Geosciences* 14, 1-13, 2021
- [9] I.E. Abraham, A.O. Yusuf, O. Oghenerume, R.A. Adelana and P.O. Azikiwe, Barite as an industrial mineral in Nigeria: occurrence, utilization, challenges and future prospects, *Heliyon*, **7(6)**, 2021.
- [10] A.O. Akanwa and N.N. Joe-Ikechebelu, Chapter 26 - Sustainable natural resources exploitation: Clay/sand mining on diminishing greener security and increased climate risks in Nigeria, *Natural Resources Conservation and Advances for Sustainability, Elsevier*, 545-562, 2022.
- [11] Geographic coordinates of Umuahia, Anonymous, Geographic coordinates of Umuahia south.
<https://www.google.com/search?q=Geographic+coordinates+of+Umuahia+south&oq=Geographic+coordinates+of+Umuahia+south&aqs=chrome..69i57j33i160l4.8089j0j15&sourceid=chrome&ie=UTF-8#bsht=Cgdic2h3Y2hwEgQIBDAB> [Assessed: 29-04-2023], [No date].
- [12] Umuahia, Anonymous, Umuahia". Wikipedia <https://en.wikipedia.org/wiki/Umuahia> [Assessed: 24-06-2023], No date.
- [13] A. Faanu, Assessment of public exposure to naturally occurring radioactive materials from mining and mineral processing activities of tarkwa gold mine in Ghana. PhD Thesis, College of Science, Kwame Nkrumah University of Science and Technology, Kumasi Ghana, 2011.
- [14] International Atomic Energy Agency, Measurement of radionuclides in food and the environment. International Atomic Energy Agency, Technical Reports Series no 295, 1989.
- [15] International Atomic Energy Agency, Extent of environmental contamination by naturally occurring radioactive material [NORM], IAEA Technical Report Series No. 419, Vienna-Austria, 2003.
- [16] International Atomic Energy Agency, Reference sheet, reference material, 2001.
- [17] G. Gilmore and J.D. Hemingway, Practical gamma spectrometry, John Wiley and Sons, NY., 1995.
- [18] T.A. Adagunodo, A.I. George, I.A. Ojoawo, K. Ojesanmi, and R. Ravisankar, Radioactivity and radiological hazards from a kaolin mining field in Ifonyintedo, Nigeria, *Science Direct Methods X journal Elsevier*, **5**:362-374, 2018.
- [19] A.A. Ibrahim, A.K. Hashim and A.A. Abojasim, Comparing of the natural radioactivity in soil samples of University at Al-Husseineya and Al-Mothafeen sites of Karbala, Iraq. *Jordan Journal of Physics*, **14(2)**, 177-191, 2021.
- [20] F. Alshahri, A. El-Taher., Investigation of Natural Radioactivity Levels and Evaluation of Radiation Hazards in Residential-Area Soil Near a Ras Tanura refinery, Saudi Arabia. *Polish Journal of Environmental Studies* 28 (1). 2019
- [21] M.O. Isinkaye, N. N. Jibiri, S.I. Bamidele and L.A. Najam, Evaluation of radiological hazards due to natural radioactivity in bituminous soils from tar-sand belt of southwest Nigeria using HpGe-Detector, *International Journal of Radiation Research*, **16(3)**, 351-362, 2018.
- [22] O. I. Araromi, A.O. Ojo, M. O. Olaluwoye and O. B. Odefemi, The Concentration of Natural Radionuclides in Soil Samples from The Practical Year Agricultural Farmland, University of Ibadan. *IOSR Journal of Applied Physics*, **8(4)**, 60-68, 2016.
- [23] M. R. Usikalu, P. P. Maleka, N. B. Ndlovu, S. Zongo, J. A. Achuka and T. J. Abodunrin, Radiation dose assessment of soil from Ijero Ekiti, Nigeria. *Cogent Engineering*, **6**, 1-11, 2019.
- [24] K.M. Ajekiigbe, F.S. Olise, S.T. Gbenu, Yinusa, S.T.V.N. Amadi and H.B. Olaniyi, Gamma Spectrometric Analysis of Soil, Sediment and Water Samples of Granitic-Type Solid Mineral Mining Activities. *Journal of Radiation and Nuclear Applications*, **2(1)**, 29-36, 2017.
- [25] United Nations Scientific Committee on the Effects of Atomic Radiation, UNSCEAR 2008, Sources and Effects of Ionizing Radiation, in: R. United Nations Scientific Committee on the Effects of Atomic [Ed.], United Nation Publication, New York, 2010.
- [26] S. P. Rihs, T. Jonathan, L. Bruno, P. Damient, C.F. Marie-Claire and F. Chabaux, using short-lived nuclides of the U-and Th-series to probe the kinetics of colloid migration in forested soils, *Geochimica et Cosmochimica Acta*, **75**, 7707-7724, 2011.
- [27] M. O. Oviru, O. U. Fredrick and A. Ochuko, Environmental risk assessment of background radiation, natural radioactivity and toxic elements in rocks and soils of Nkalagu quarry, Southeastern Nigeria, *Journal of Hazardous Materials Advances*, **10**, 2023.
- [28] T.O. Kolawole, O. Iyiola, H. Ibrahim and R. A. Isibor, Contamination, ecological and health risk assessments of potentially toxic elements in soil around a municipal solid waste disposal facility in Southwestern Nigeria, *Journal of Trace Elements and Minerals*, **5**, 100083, 2023.
- [29] A.J. Innocent, M.Y. Onimisi, and S.A. Jonah, Evaluation of naturally occurring radionuclide materials

- in soil samples collected from some mining sites in Zamfara State, Nigeria, *British Journal of Applied Science and Technology*, **3(4)**, 684-692, 2013.
- [30] HMH Zakaly, MAM Uosif, SAM Issa, HO Tekin, H Madkour, M Tammam, ...An extended assessment of natural radioactivity in the sediments of the mid-region of the Egyptian Red Sea coast. *Marine Pollution Bulletin* 171, 112658. 2021.
- [31] J.C. Osimobi, G.O. Avwiri, and E.O. Agbalagba, Radiometric and radiogenic heat evaluation of natural radioactivity of soil around solid minerals mining environment in south-eastern Nigeria. *Environmental Processes an International Journal, Springer Nature*, **5(4)**, 859-877, 2018.
- [32] J.A. Santos Júnior, E.E.N. Araújo, Z.H. Fernández, R.S. Amaral, J.M.N. Santos, M.O. Milán, Measurement of natural radioactivity and radium equivalent activity for pottery making clay samples in Paraíba and Rio Grande do Norte – Brazil, *Environmental Advances*, **6**, 100121, 2021.
- [33] U.K. Mayeen, A. Khandoker, F.B.S. Abdullah, D.A. Bradley and M.O. Isinkay, Elevated concentrations of naturally occurring radionuclides in heavy mineral-rich beach sands of Langkawi Island, Malaysia, *Marine Pollution Bulletin*, **127**, 654-663, 2018.
- [34] A. El-TaHER, HA Madkour., Environmental and radioecological studies on shallow marine sediments from harbour areas along the Red Sea coast of Egypt for identification of anthropogenic impacts. *Isotopes in environmental and health studies* 50 (1), 120-133. 2014.
- [35] WM Badawy, A El-TaHER, MV Frontasyeva, HA Madkour, AEM Khater., Assessment of anthropogenic and geogenic impacts on marine sediments along the coastal areas of Egyptian Red Sea. *Applied Radiation and Isotopes* 140, 314-326. 2018.
- [36] D.A. Fungaro, P.S.C. Silva, F.A. Campello, C.S. Miranda and J. C. Izidoro, Evaluation of radionuclide contamination of soil, coal ash and zeolitic materials from Figueira Thermoelectric Power Plant, *Brazilian Journal of Radiation Sciences*, 01-18, 2019.
- [37] S. Singh, A. Rani, and R.K. Mahajan, ^{226}Ra , ^{232}Th and ^{40}K analysis in soil samples from some areas of Punjab and Himachal Pradesh, India using gamma ray spectrometry, *Radiation Measurements*, **39(4)**, 431-439, 20005.
- [38] S. Dizman, F.K. Görür, R. Keser, Determina on of radioactivity levels of soil samples and the excess of life me cancer risk in Rize province, Turkey. *International Journal of Radiation Research*, **14(3)**, 237-244, 2016.
- [39] A.J. Innocent, M.Y. Onimisi and Jonah, S.A, Evaluation of naturally occurring radionuclide materials in soil samples collected from some mining sites in Zamfara State, Nigeria, *British Journal of Applied Science and Technology*, **3(4)**, 684, 2013.
- [40] E.O. Agbalagba and R.A. Onoja, Evaluaon of natural radioacvity in soil, sediment and water samples of Niger Delta Biseni flood plain lakes, Nigeria. *Journal of Environmental Radioacvity*, **102(7)** 667-671, 2011.
- [41] O. A. Oyebanjo, E. O. Falayi, F. O. Ogunsanwo and A. I. Odudu, Natural radioactivity and hazards in soil samples in Alade community, Southwestern Nigeria, *Journal of Physical and Life Sciences*, **1(21)**, 2018.
- [42] A. A. Mbet, U. Ibrahim and I. Shekwonyadu, Assessment of radiological risk from the soils of artisanal mining areas of Anka, North West Nigeria, African, *Journal of Environmental Science and Technology*, **3(8)**, 303-309, 2019.
- [43] S. Fares, Measurements of natural radioactivity level in black sand and sediment samples of the Tamsah Lake beach in Suez Canal region in Egypt, *Journal of Radiation Research and Applied Sciences*, **10(3)**, 194-203, 2017.
- [44] E. Devanesan, J. Chandramohan, G. Senthilkumar, N. Harikrishnan, M.S. Gandhi, S.S. Kolekar and R. Ravisankar, Natural radioactivity concentrations and dose assessment in coastal sediments along the East Coast of Tamilnadu, India with statistical approach, *Acta Ecologica Sinica*, **40(5)**, 353-362, 2020.
- [45] G., Akkaya, G., Kaynak, A., Kahraman and O. Gurler, The investigation of radionuclide distributions in soil samples collected from Bursa, Turkey, *Radiation Protection Dosimetry*, **152(4)**, 376-383, 2017.
- [46] A. Y. Ahmad, M. A. Al-Ghouti, I. AlSadig N. and M. Abu-Dieyeh, Vertical distribution and radiological risk assessment of ^{137}Cs and natural radionuclides in soil samples, *Sci Rep*, **9(1)**, 12196 2019.